
State of the Bay

Casco Bay Estuary Partnership (CBEP)

2010

Interior Forest Habitat (2010 State of the Bay Poster)

Casco Bay Estuary Partnership

Follow this and additional works at: <https://digitalcommons.usm.maine.edu/cbep-state-of-the-bay>

Recommended Citation

Casco Bay Estuary Partnership. (2010). Interior Forest Habitat (2010 State of the Bay Poster). [Poster]. Portland, ME: University of Southern Maine, Muskie School of Public Service, Casco Bay Estuary Partnership.

This Book is brought to you for free and open access by the Casco Bay Estuary Partnership (CBEP) at USM Digital Commons. It has been accepted for inclusion in State of the Bay by an authorized administrator of USM Digital Commons. For more information, please contact jessica.c.hovey@maine.edu.

Interior Forest Habitat



Peter Taylor/WaterView Consulting


Forests provide essential habitat to many of Maine's native birds, fish, and mammals. Certain species, including large herbivores and predators such as fishers, hawks and owls, roam over large areas of forest and thus cannot survive in the small forests found in suburban areas. Many species of migrant songbirds, including many warblers, are forest specialists, nesting successfully only in large blocks of forest.

While to most humans there may appear to be little difference between the edge and the interior of a forest, there can be profound differences from the perspective of the animals and plants that live there.

Even in a largely forested watershed such as Casco Bay, suitable habitat for forest specialists may be uncommon. Their ideal habitat occurs only in large areas of forest that are compact in shape and are located far from most human activity.

- Almost 69 percent (676.0 square miles) of the 986 square mile Casco Bay watershed is forested (Maine Office of GIS 2004).
- In contrast, only 172.6 square miles (17.5 percent) of the watershed consists of interior forest habitat, the majority of which is located in the upper portions of the watershed.
- Interior forest is far less abundant in the more highly developed coastal communities, where suburban lands, abundant roads, powerlines, and other linear infrastructure cut the forest into smaller areas that provide little true interior forest habitat.

Interior Forest Habitat in Casco Bay Townships



The majority of the interior forest habitat in the Casco Bay watershed lies within the northern and western towns at the headwaters of the Sebago Lake / Presumpscot River watershed. The more developed coastal communities contain little or no interior forest habitat.

