

2013

Clam Abundance and PH: Preliminary Data From Casco Bay (2013 Mud Summit Presentation)

Curtis C. Bohlen PhD
Casco Bay Estuary Partnership

Follow this and additional works at: <https://digitalcommons.usm.maine.edu/cbep-presentations>

Recommended Citation

Bohlen, C. C.. (2013). Clam Abundance and PH: Preliminary Data From Casco Bay (2013 Mud Summit Presentation). [Presentation slides]. Portland, ME: University of Southern Maine, Muskie School of Public Service, Casco Bay Estuary Partnership.

This Book is brought to you for free and open access by the Casco Bay Estuary Partnership (CBEP) at USM Digital Commons. It has been accepted for inclusion in Presentations by an authorized administrator of USM Digital Commons. For more information, please contact jessica.c.hovey@maine.edu.

CLAM ABUNDANCE AND PH – PRELIMINARY DATA FROM CASCO BAY

Mud Summit, January 18, 2013


Curtis C. Bohlen
Director, Casco Bay Estuary Partnership



Special Thanks


- The people who did the field work
 - Chris Heinig (MER Assessment)
 - Steve Karpiak (MER Assessment)
 - Matthew Craig
 - Caitlin Gerber
 - Cayce Dalton (FB Environmental)

Clam Flat pH Sites, 2011




Site Selection (Reminder)

- Sites selected by contacting local informants in each town
- Identify productive flats and flats that are no longer considered productive




pH and Informant Assessment of Productivity



- “Productive” flats have higher average pH than “Unproductive” Flats
- Difference =
 0.23 ± 0.101
($p < 0.05$)

Methods

- Clam abundance data collected in November
- Return to recorded GPS position of FOCB pH measurement
- Eight spat and eight clam samples collected radially around that point
 - ▣ 10cm diameter PVC pipe Spat cores
 - ▣ 0.1 m² rectangular Clam sampling plot (hand dug)




Productivity of Flats

Informant Assessment of Productivity				
	Unproductive	Productive	Total	
Observed Productivity	Unproductive	13	9	22
	Productive	2	6	8
	Total	15	15	30

- When researchers visited more than one third of the “Productive” flats, they found few clams near the sites where pH had been measured.
- But in some cases, clams were abundant nearby.

Adults and Spat are Correlated

- Six flats had spat, but no adults
- Three flats had adults, but no spat
- Spat and adult abundance are correlated, but not strongly



Preliminary Conclusions

- Data on relation between pH and shellfish abundance is suggestive but not conclusive
- Sample locations in many flats were located in areas with few clams, reducing sensitivity
- Better understanding of spatial pattern is needed
 - ▣ Spatial distribution of clams
 - ▣ Spatial autocorrelation function of pH
- Future studies need
 - ▣ Improved sampling design
 - ▣ More complete measurement of environmental correlates

